REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

2. REPORT TYPE

3. DATES COVERED (From - To)

28 March 2003	View Graphs			
4. TITLE AND SUBTITLE			58	. CONTRACT NUMBER
New Polynitrogen Compounds			5t	O. GRANT NUMBER
			50	:. PROGRAM ELEMENT NUMBER
6. AUTHOR(S) Jerry A. Boatz, Karl Christie, Ralph Ha	aiges, Stefan Schneider,	Thorsten Schroer, As	I	I. PROJECT NUMBER 303
Vij, William W. Wilson				e. TASK NUMBER 12C8
			5f	. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			PERFORMING ORGANIZATION REPORT UMBER
Air Force Research Laboratory (AFMO AFRL/PRSP & ERC., Inc.	2)		A	FRL-PR-ED-VG-2003-076
10 E. Saturn Blvd. Edwards AFB, CA 93524-7680				
9. SPONSORING / MONITORING AGENCY		55(ES)	10). SPONSOR/MONITOR'S ACRONYM(S)
Air Force Research Laboratory (AFMO	2)			
AFRL/PRS		•	11	. SPONSOR/MONITOR'S
5 Pollux Drive				NUMBER(S) FRL-PR-ED-VG-2003-076
Edwards AFB CA 93524-7048			A	FRL-FR-ED-VG-2003-070
12. DISTRIBUTION / AVAILABILITY STAT	EMENT			
Approved for public release; distribution	on unlimited.			
13. SUPPLEMENTARY NOTES				
14. ABSTRACT				
			0.00	30000
			200	30805 153
				70007 175
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	PERSON
	- TINO DAGE	_		Sheila Benner
a. REPORT b. ABSTRACT	c. THIS PAGE	A		19b. TELEPHONE NUMBER (include area code) (661) 275-5963
Unclassified Unclassified	Unclassified		<u> </u>	Standard Form 298 (Rev. 8-98)
				Prescribed by ANSI Std. 239.18



MEMORANDUM FOR PRS (In-House Publication)

FROM: PROI (STINFO)

31 Mar 2003

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-VG-2003-076 Karl Christe (AFRL/PRSP (ERC)); Ralf Haiges, Stefan Schneider, Thorsten Schroer (Loker Hydrocarbon Research Institute); Ashwani Vij, Vandana Vij, and William W. Wilson (AFRL/PRSP (ERC)), "New Polynitrogen Compounds"


AFOSR Molecular Dynamics Contractors Conference (San Diego, CA, 18-20 May 2003) (Deadline: 18 May 2003)

(Statement A)

5364 Christe 5194

Towards New Polynitrogen Species: The Search for Viable Precursors **AFOSR Molecular Dynamics Contractors**

Conference May 18-20, 2003



Dr. Jerry A. Boatz
Senior Research Chemist
Propellants Branch
Air Force Research Laboratory
jerry.boatz@edwards.af.mil
(661) 275-5364

DISTRIBUTION STATEMENT AApproved for Public Release
Distribution Unlimited



HEDM Program Objective







- Hydrocarbons for liquid boosters
- Liquid & solid oxidizers for boost and upper stages
- · Monopropellants for upper stages and satellites











Current State of the Art



Monopropellants	Isp (sec)	lsp (sec) Density (g/cm³)
Hydrazine (N ₂ H ₄)	200	1.01
Hydrogen Peroxide (H ₂ O ₂)	175	1.41
Solids		
Ammonium Perchlorate (AP)	260	1.69
~20%AI/~10% polymer binder		
Liquids		
N ₂ O₄/Hydrazine	280	1.45/1.01
RP-1/Liquid Oxygen (LOX)	300	0.81/1.15
Cryogenic		
Liquid Hydrogen (LH2)/LOX	390	0.07/1.15
I _{sp} at sea level 1000 psi chamber pressure	psi chamber p	ressure
lsp = (pounds of thrust/pounds of propellant/second)	unds of propell	ant/second)



HEDM Propellant Payoffs



"The highest leverage technology area impacting launch vehicles is the development of high-energy-density materials for use as propellants." -- New World Vistas Panel on Space Technology (1995)

Propellant Takeoff Mass (lb)
RP-1/LOX 360,000
(lsp = 295 s) // LH2/LOX
(lsp = 455 s)
LH2/LOX 1,900,000
(1sp = 455 s)
HTPB/AI/HMX 1,847
(lsp = 270 s)

Our research is aimed at increasing propellant Isp by 5 to 50%




Polynitrogen Project




Discover, synthesize, characterize, and scale-up novel, highly energetic polynitrogen allotropes


Modeling and simulation guides the experimental program:

- Determines which molecules should exist and how energetic they are
- Gives information on how to synthesize promising molecules
- Provides critical data for identification and characterization of new molecules
















Polynitrogen Monopropellants Performance of



The performance of polynitrogens as monopropellants would dwarf that of hydrazine, and would greatly exceed even bipropellants





The Search for New Polynitrogens



- All polynitrogens are unstable with respect to N₂ molecules
- Their activation energy for N₂ elimination is largely determined by the weakest bond in the compound
- Their metastability is enhanced if suitable resonance structures exist:

$$-[|\overline{N}-N=N|] \longleftrightarrow -[\sqrt{N=N-N}] \longleftrightarrow -[|N=N-\overline{N}|]$$

- The double-bond character of the N-N bonds in the azide anion explains its exceptional stability
- How can this stabilization effect be used to our advantage in preparing new compounds?



Pentazolate (N₅-)?




- Substituted pentazoles R-N₅ have been known for decades (R=aryl)
- Cyclic N₅- is aromatic
- Conversion of the diazonium salt, RN₂⁺, to the substituted pentazole ring R-N₅ by the reaction with azide ion, N_3 -, has been demonstrated many years ago by Ugi and Huisgen.
- N₅- has been recently detected in the gas phase for the first time, using collisional fragmentation (electrospray ion mass spectroscopy).
- Can a chemical route to N₅- be found? (e.g., can a suitable R group be found for the reaction R-N₂⁺ + N₃⁻ -> R-N₅ -> R⁺ + N₅⁻?




New Polynitrogen Anions





Pentazole anion (N₅-)

- Theoretical calculations show that this anion has a 28 kcal/mole activation energy barrier for decomposition and its decomposition to N_3^- and N_2 is only 11 kcal/mol exothermic
- Aryl substituted pentazoles can be isolated as stable compounds only if stored at low temperatures. In methanol, these compounds rapidly decompose at room temperature to form aryl azides and N₂ gas





Synthetic Challenge - How do we make These New Anions??




Synthesis of Substituted Pentazoles

Sources for the Pentazole Anion (N₅-)



Aryl Diazonium Salts





Theoretical Challenge - Can we design and predict viable precursors to N₅-?



Find a substituent R so that

- 1. R-N₂⁺ is stable wrt R⁺ + N₂.
- 2. The R-N bond in R-N₅ is weak, thereby suitable for the reaction R-N₅ + M⁺X⁻ -> M⁺N₅ + R-X

Approach

stability of R-N₂⁺ diazonium salts and the length/strength Use quantum chemical calculations to predict the of the R-N₅ bond.

MBPT(2)/6-31G(d) level of theory

$$L_3M-N_2^+$$


$$M = C, Si$$


$$L = F$$
, CI, CH₃, CF₃, NO₂, phenyl, etc.



Phenyldiazonium -- too stable!







 $R(C-N_5) = 1.43 \text{ Å}$

$$R(C-N_2^+) = 1.39 \text{ Å}$$


$$D_e = 37.9 \text{ kcal/mol}$$


Try replacing Ph+ with a more stable cation (Ph₃C⁺)...



Trityldiazonium -- unstable!








placing e-withdrawing groups on the phenyl rings (e.g. NO₂)? Trityl cation Ph₃C+ is too stable -- can it be destabilized by



tris(para-nitrophenyl) methyldiazonium





R₃C-N₂⁺ distance(Å) D_e(C-N₂⁺) (kcal/mol)

> 3.0

> 3.0

 $3(pNO_2-Ph)$


3Ph




What about some silicon analogues?




D _e (Si-N ₂ ⁺) (kcal/mol)	-6.0	-13.8
R ₃ Si-N ₂ ⁺ distance(Å)	> 2.5	2.151
R _s	3Ph	ЗМе













Derivatives of trimethylsilyldiazonium



$se(\text{Å}) D_e(Si-N_2^+) \text{ (kcal/mol)}$	-7.4	-5.2
R ₃ Si-N ₂ ⁺ distance(Å)	2.410	> 2.5
R ₃ R ₃ Si	2Me,FCH ₂	2Me,NO ₂ CH ₂











Silicon analogues, continued

R 3	R ₃ Si-N ₂ ⁺ distance(Å)	D _e (Si-N ₂ ⁺) (kcal/mol)
3F	1.973	-33.4
3CI	2.125	-14.5
3CF ₃	1.994	n/a













Conclusions



- Trityldiazonium (Ph₃C-N₂⁺) and its tris(para-NO₂) analogue are predicted to be unstable wrt dissociation of N₂
- Ph₃Si-N₂⁺ is also predicted to be unstable wrt dissociation of N₂
- Me₃Si-N₂⁺ and Cl₃Si-N₂⁺ are marginally stable (D_e $\sim 14 \text{ kcal/mol}$; R(Si-N) $\sim 2.1 \text{ Å}$)
- Of the silyl derivatives considered, $F_3Si-N_2^+$ is the most stable ($D_e=33.4$ kcal/mol; $R(Si-N_2^+) = 1.97 \text{ Å})$
- NO₂ interact strongly with Si in silyldiazonium cations, leading to displacement of Secondary (i.e., not directly bonded to Si) electronegative groups such as F and



Future/ongoing work



 Calculation of the structures and stabilities of Me_xF_yCl_zSi-N₂⁺ and $Me_xF_yCI_zSi-N_5$ • Calculation of the reaction enthalpies R-N₅ + M^+X^- -> $M^+N_5^-$ + R-X

Other high-nitrogen/polynitrogen species (e.g. Fe(N₅)₂)





DARPA, AFOSR, AFRL

DoD HPC Modernization Program

New Materials Design Challenge Project

Aeronautical System Center

AFFTC Distributed Center

Army HPC Research Center

Common HPC Software Support Initiative

Dr. Robert Corley (AFRL)

